

# BRAIN RESEARCH

## MOLECULAR BRAIN RESEARCH

VOL. 46 NOS. 1,2

### CONTENTS

JUNE 1997

Cited in Biological Abstracts (BIOSIS) – Chem. Abstracts – Index Medicus (MEDLINE) – Current Contents (Life Sci.) – Excerpta Medica (EMBASE) – Psychological Abstracts (PsycINFO) – Pascal et Francis (INIST-CNRS) – RIS (Reference Update) – Current Awareness in Biological Science (CABS)

Publisher's Note

v

Guidelines for the submission of manuscripts

vii

Research reports

|   |     |
|---|-----|
| Opioid receptor expression in the rat gastrointestinal tract: a quantitative study with comparison to the brain<br>J. Fickel, D. Bagnol, S.J. Watson and H. Akil (USA)  | 1   |
| Differential proopiomelanocortin gene expression in the medial basal hypothalamus of rats during pregnancy and lactation<br>P.E. Mann, B.S. Rubin and R.S. Bridges (USA)  | 9   |
| Extensive splice variation and localization of the EHK-1 receptor tyrosine kinase in adult human brain and glial tumors<br>G.C. Miescher, V. Taylor, G. Olivieri, T. Mindermann, E. Schröck and A.J. Steck (Switzerland, Germany)                             | 17  |
| Chronic food restriction and streptozotocin-induced diabetes differentially alter prodynorphin mRNA levels in rat brain regions<br>Y. Berman, L. Devi, R. Spangler, M.J. Kreek and K.D. Carr (USA)  | 25  |
| Deprenyl induces the tyrosine hydroxylase enzyme in the rat dopaminergic nigrostriatal system<br>J.A. Rodríguez-Gómez, J.L. Venero, M.L. Vizcute, J. Cano and A. Machado (Spain)  | 31  |
| Aging-related regulation of myo-inositol 1,4,5-trisphosphate signal transduction pathway in the rat striatum<br>O.J. Igwe and M.B. Filla (USA)  | 39  |
| Regional and cellular expression patterns of four K <sup>+</sup> channel mRNAs in the adult rat brain<br>S. Verma-Kurvari, B. Border and R.H. Joho (USA)  | 54  |
| Olfactory memory and maternal behaviour-induced changes in c-fos and zif/268 mRNA expression in the sheep brain<br>A.P.C. Da Costa, K.D. Broad and K.M. Kendrick (UK)   | 63  |
| Effects of ethanol on muscarinic receptor-stimulated c-fos expression in human neuroblastoma cells<br>W.-Q. Ding, C. Larsson, P. Simonsson and C. Alling (Sweden)   | 77  |
| An anti- $\chi$ -1 antibody recognizes a heavily glycosylated protein in rat brain<br>A.M. Ciabarra and K.A. Sevarino (USA)   | 85  |
| Transgene expression of plasmid DNAs directed by viral or neural promoters in the rat brain<br>Z. Hannas-Djebara, M. Didier-Bazès, S. Sacchettini, C. Prod'hon, M.-F. Belin and B. Jacquemont (France)  | 91  |
| Effects of chronic exposure to $\Delta^9$ -tetrahydrocannabinol on cannabinoid receptor binding and mRNA levels in several rat brain regions<br>J. Romero, E. García-Palomero, J.G. Castro, L. García-Gil, J.A. Ramos and J.J. Fernández-Ruiz (Spain)         | 100 |
| Expression of pituitary adenylate cyclase-activating polypeptide (PACAP) in the mesencephalic trigeminal nucleus of the rat after transection of the masseteric nerve<br>J. Oemar Larsen, J. Hannibal, S.M. Knudsen and J. Fahrenkrug (Denmark)               | 109 |
| Nerve growth factor mRNA stability is controlled by a cis-acting instability determinant in the 3'-untranslated region<br>B. Tang, M. Wang and B.C. Wise (USA)  | 118 |
| Comparison of the effects of an ampakine with those of methamphetamine on aggregate neuronal activity in cortex versus striatum<br>L.C. Palmer, U.S. Hess, J. Larson, G.A. Rogers, C.M. Gall and G. Lynch (USA)   | 127 |
| Expression and functional role of the Id HLH family in cultured astrocytes<br>S.-F. Tzeng and J. De Velis (USA)   | 136 |
| NGFI-A expression in the rat brain after sleep deprivation<br>M. Pompeiano, C. Cirelli, S. Ronca-Testoni and G. Tononi (Italy)  | 143 |
| Elevated expression of glutathione peroxidase in PC12 cells results in protection against methamphetamine but not MPTP toxicity<br>D.G. Hom, D. Jiang, E.-J. Hong, J.Q. Mo and J.K. Andersen (USA)  | 154 |
| Pharmacological modulation of Alzheimer's $\beta$ -amyloid precursor protein levels in the CSF of rats with forebrain cholinergic system lesions<br>V. Haroutunian, N. Greig, X.-F. Pei, T. Utsuki, R. Gluck, L.D. Acevedo, K.L. Davis and W.C. Wallace (USA) | 161 |
| Distribution and polysome association of full-length and 3' truncated preproenkephalin mRNA in the guinea-pig brain<br>K.S. LaForge, A. Ho and M.J. Kreek (USA)   | 169 |
| Induction of FOS and JUN proteins during focal epilepsy: congruences with and differences to [ <sup>14</sup> C]deoxyglucose metabolism<br>P. Gass, C. Bruehl, T. Herdegen, M. Kiessling, M. Lutzenburg and O.W. Witte (Germany)                               | 177 |
| Cloning and characterization of Kir3.1 (GIRK1) C-terminal alternative splice variants<br>C.S. Nelson, J.L. Marino and C.N. Allen (USA)  | 185 |

(contents continued on p. 350)



0169-328X(199706)46:1/2;1-0

# BRAIN RESEARCH

## MOLECULAR BRAIN RESEARCH

VOL. 46 NOS. 1,2

### CONTENTS

JUNE 1997

Cited in Biological Abstracts (BIOSIS) – Chem. Abstracts – Index Medicus (MEDLINE) – Current Contents (Life Sci.) – Excerpta Medica (EMBASE) – Psychological Abstracts (PsycINFO) – Pascal et Francis (INIST-CNRS) – RIS (Reference Update) – Current Awareness in Biological Science (CABS)

Publisher's Note

v

Guidelines for the submission of manuscripts

vii

Research reports

|   |     |
|---|-----|
| Opioid receptor expression in the rat gastrointestinal tract: a quantitative study with comparison to the brain<br>J. Fickel, D. Bagnol, S.J. Watson and H. Akil (USA)  | 1   |
| Differential proopiomelanocortin gene expression in the medial basal hypothalamus of rats during pregnancy and lactation<br>P.E. Mann, B.S. Rubin and R.S. Bridges (USA)  | 9   |
| Extensive splice variation and localization of the EHK-1 receptor tyrosine kinase in adult human brain and glial tumors<br>G.C. Miescher, V. Taylor, G. Olivieri, T. Mindermann, E. Schröck and A.J. Steck (Switzerland, Germany)                             | 17  |
| Chronic food restriction and streptozotocin-induced diabetes differentially alter prodynorphin mRNA levels in rat brain regions<br>Y. Berman, L. Devi, R. Spangler, M.J. Kreek and K.D. Carr (USA)  | 25  |
| Deprenyl induces the tyrosine hydroxylase enzyme in the rat dopaminergic nigrostriatal system<br>J.A. Rodríguez-Gómez, J.L. Venero, M.L. Vizcute, J. Cano and A. Machado (Spain)  | 31  |
| Aging-related regulation of myo-inositol 1,4,5-trisphosphate signal transduction pathway in the rat striatum<br>O.J. Igwe and M.B. Filla (USA)  | 39  |
| Regional and cellular expression patterns of four K <sup>+</sup> channel mRNAs in the adult rat brain<br>S. Verma-Kurvari, B. Border and R.H. Joho (USA)  | 54  |
| Olfactory memory and maternal behaviour-induced changes in c-fos and zif/268 mRNA expression in the sheep brain<br>A.P.C. Da Costa, K.D. Broad and K.M. Kendrick (UK)   | 63  |
| Effects of ethanol on muscarinic receptor-stimulated c-fos expression in human neuroblastoma cells<br>W.-Q. Ding, C. Larsson, P. Simonsson and C. Alling (Sweden)   | 77  |
| An anti- $\chi$ -1 antibody recognizes a heavily glycosylated protein in rat brain<br>A.M. Ciabarra and K.A. Sevarino (USA)   | 85  |
| Transgene expression of plasmid DNAs directed by viral or neural promoters in the rat brain<br>Z. Hannas-Djebara, M. Didier-Bazès, S. Sacchettini, C. Prod'hon, M.-F. Belin and B. Jacquemont (France)  | 91  |
| Effects of chronic exposure to $\Delta^9$ -tetrahydrocannabinol on cannabinoid receptor binding and mRNA levels in several rat brain regions<br>J. Romero, E. Garcia-Palomero, J.G. Castro, L. Garcia-Gil, J.A. Ramos and J.J. Fernandez-Ruiz (Spain)         | 100 |
| Expression of pituitary adenylate cyclase-activating polypeptide (PACAP) in the mesencephalic trigeminal nucleus of the rat after transection of the masseteric nerve<br>J. Oemar Larsen, J. Hannibal, S.M. Knudsen and J. Fahrenkrug (Denmark)               | 109 |
| Nerve growth factor mRNA stability is controlled by a cis-acting instability determinant in the 3'-untranslated region<br>B. Tang, M. Wang and B.C. Wise (USA)  | 118 |
| Comparison of the effects of an ampakine with those of methamphetamine on aggregate neuronal activity in cortex versus striatum<br>L.C. Palmer, U.S. Hess, J. Larson, G.A. Rogers, C.M. Gall and G. Lynch (USA)   | 127 |
| Expression and functional role of the Id HLH family in cultured astrocytes<br>S.-F. Tzeng and J. De Velis (USA)   | 136 |
| NGFI-A expression in the rat brain after sleep deprivation<br>M. Pompeiano, C. Cirelli, S. Ronca-Testoni and G. Tononi (Italy)  | 143 |
| Elevated expression of glutathione peroxidase in PC12 cells results in protection against methamphetamine but not MPTP toxicity<br>D.G. Hom, D. Jiang, E.-J. Hong, J.Q. Mo and J.K. Andersen (USA)  | 154 |
| Pharmacological modulation of Alzheimer's $\beta$ -amyloid precursor protein levels in the CSF of rats with forebrain cholinergic system lesions<br>V. Haroutunian, N. Greig, X.-F. Pei, T. Utsuki, R. Gluck, L.D. Acevedo, K.L. Davis and W.C. Wallace (USA) | 161 |
| Distribution and polysome association of full-length and 3' truncated preproenkephalin mRNA in the guinea-pig brain<br>K.S. LaForge, A. Ho and M.J. Kreek (USA)   | 169 |
| Induction of FOS and JUN proteins during focal epilepsy: congruences with and differences to [ <sup>14</sup> C]deoxyglucose metabolism<br>P. Gass, C. Bruehl, T. Herdegen, M. Kiessling, M. Lutzenburg and O.W. Witte (Germany)                               | 177 |
| Cloning and characterization of Kir3.1 (GIRK1) C-terminal alternative splice variants<br>C.S. Nelson, J.L. Marino and C.N. Allen (USA)  | 185 |

(contents continued on p. 350)



0169-328X(199706)46:1/2;1-0

## (contents continued)

|   |     |
|---|-----|
| Tissue-specific methylation occurs in the essential promoter element of the tyrosine hydroxylase gene<br>K. Okuse, I. Matsuoaka and K. Kurihara (Japan)   | 197 |
| Transcriptional regulation of the human S100 $\beta$ gene<br>F. Castets, W.S.T. Griffin, A. Marks and L.J. Van Eldik (USA, Canada)  | 208 |
| Localization of mRNA for fatty acid transport protein in developing and mature brain of rats<br>A. Utsunomiya, Y. Owada, T. Yoshimoto and H. Kondo (Japan)  | 217 |
| Distribution of the neuropeptide Y Y2 receptor mRNA in rat central nervous system<br>E.L. Gustafson, K.E. Smith, M.M. Durkin, M.W. Walker, C. Gerald, R. Weinstock and T.A. Brancheck (USA)   | 223 |
| Induction of Na $^+$ /myo-inositol co-transporter mRNA after rat cryogenic injury<br>T. Yamashita, S. Shimada, A. Yamauchi, W. Guo, E. Kohmura, T. Hayakawa and M. Tohyama (Japan)  | 236 |
| The melanin-concentrating hormone gene in human: flanking region analysis, fine chromosome mapping, and tissue-specific expression<br>A. Viale, Y. Zhixing, C. Breton, F. Pedeutour, A. Coquerel, D. Jordan and J.-L. Nahon (France)  | 243 |
| Differential distribution of mRNAs encoding phosphatidylinositol transfer proteins $\alpha$ and $\beta$ in the central nervous system of the rat<br>H. Imai, S. Tanaka, T. Fukusato, S. Yamashita and K. Hosaka (Japan)   | 256 |
| A newly identified membrane protein localized exclusively in intracellular organelles of neurons<br>Y. Kadota, A. Niiya, R. Masaki, A. Yamamoto, M. Araki and S. Taketani (Japan)   | 265 |
| TrkB expression in dentate granule cells is associated with a late phase of long-term potentiation<br>M. Dragunow, P. Hughes, S.E. Mason-Parker, P. Lawlor and W.C. Abraham (New Zealand, USA)  | 274 |
| Distribution of AMPA receptor subunit mRNAs in the human basal ganglia: an in situ hybridization study<br>M. Tomiyama, J.M. Palacios, R. Cortés, M.T. Vilaró and G. Mengod (Spain)  | 281 |
| Regulation of hippocampal 5-HT <sub>1A</sub> receptor mRNA and binding in transgenic mice with a targeted disruption of the glucocorticoid receptor<br>O.C. Meijer, T.J. Cole, W. Schmid, G. Schütz, M. Joëls and E.R. De Kloet (The Netherlands, Germany)                        | 290 |
| Induction of heat-shock protein (HSP72) in the cingulate and retrosplenial cortex by drugs that antagonize the effects of excitatory amino acids<br>J.Q. Lan, J. Chen, F.R. Sharp, R.P. Simon and S.H. Graham (USA)   | 297 |
| Visual sensitivities of <i>nur77</i> (NGFI-B) and <i>zif268</i> (NGFI-A) induction in the suprachiasmatic nucleus are dissociated from <i>c-fos</i> induction and behavioral phase-shifting responses<br>J.T. Lin, J.M. Komhauser, N.P. Singh, K.E. Mayo and J.S. Takahashi (USA) | 303 |
| Identification in the rat neurotensin receptor of amino-acid residues critical for the binding of neurotensin<br>J.-M. Botto, J. Chabry, D. Nouel, M. Paquet, P. Séguéla, J.-P. Vincent, A. Beaudet and J. Mazella (France, Canada)   | 311 |
| <i>Short communications</i>   |     |
| Temporal and spatial expression of glutamic acid decarboxylases in human fetal brain<br>S.O. Chan, W.D. Lyman and F.-C. Chiu (USA)  | 318 |
| Opioidergic and dopaminergic gene expression in the caudate-putamen and accumbens of the mutant mouse, tottering ( <i>tg/tg</i> )<br>A. De Bartolomeis, V. Koprivica, D. Pickar, J.N. Crawley and L.C. Abbott (USA)   | 321 |
| Anoxia regulates gene expression in the central nervous system of <i>Drosophila melanogaster</i><br>E. Ma and G.G. Haddad (USA)   | 325 |
| Molecular cloning of a new unc-33-like cDNA from rat brain and its relation to paraneoplastic neurological syndromes<br>T.T. Quach, Y. Rong, M.F. Belin, A.-M. Duchemin, H. Akaoka, S. Ding, M. Baudry, P.E. Kolattukudy and J. Honnorat (France, USA)                            | 329 |
| Absence of neuroanatomical and behavioral deficits in L7/pcp-2-null mice<br>G. Vassileva, R.J. Smeyne and J.I. Morgan (USA)   | 333 |
| Calcium- and calmodulin-dependent phosphorylation of AMPA type glutamate receptor subunits by endogenous protein kinases in the post-synaptic density<br>Y. Hayashi, A. Ishida, H. Katagiri, M. Mishina, H. Fujisawa, T. Manabe and T. Takahashi (Japan)                          | 338 |
| Ovariectomy and estradiol treatment affect the dopamine transporter and its gene expression in the rat brain<br>R. Bossé, R. Rivest and T. Di Paolo (Canada)  | 343 |
| <i>Author Index</i>   |     |
|   | 347 |